

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
8 April 2004 (08.04.2004)


PCT

(10) International Publication Number
WO 2004/028694 A1

- (51) International Patent Classification⁷: **B02B 1/04**
- (21) International Application Number:
PCT/EP2003/010804
- (22) International Filing Date:
29 September 2003 (29.09.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
MI2002A002066
30 September 2002 (30.09.2002) IT
- (71) Applicant (*for all designated States except US*): **BAR-
ILLA ALIMENTARE S.P.A.** [IT/IT]; Via Mantova, 166,
I-43100 Parma (IT).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **TRIBUZIO, Gio-
vanni** [IT/IT]; Via Telesforo, 31, I-71100 Foggia (IT).
LODI, Alberto [IT/IT]; Via Bellini, 48, I-43023 Monticelli
(IT). **GOTTOFREDI, Angelo** [IT/IT]; Via Guareschi,
18/a, I-43036 Fidenza (IT). **RANIERI, Roberto** [IT/IT];
Via Sanguigna, 130, I-43052 Colomo (IT).
- (74) Agents: **BOTTI, Mario** et al.; Botti & Ferrari S.r.l., Via
Locatelli, 5, I-20124 Milano (IT).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE,
GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR,
KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK,
MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT,
RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
- with international search report
 - before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments
- For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: A PROCESS FOR THE PRODUCTION OF WHEAT FLOUR

(57) Abstract: A process for the production of wheat flour or semolina, comprising the steps of a) wetting the caryopses of wheat with such an amount of water as to bring their moisture content to at least 15 %, subjecting them to intense vibrations; b) subjecting the wet caryopses to a conditioning step; c) subjecting the conditioned caryopses to operations of decortication, to take off the outer layers of bran; d) milling the conditioned and decorticated caryopses.



WO 2004/028694 A1

Title: "A process for the production of wheat flour "

DESCRIPTION

The present invention relates to the foodstuff industry.

5 In particular it relates to a process for the production of wheat flour and especially semolina from durum wheat (*Triticum turgidum* var. *durum*, Desf.).

As is known, flour and semolina, respectively from soft wheat (*Triticum aestivum*, L.) and durum wheat, are obtained by grinding the caryopses and subsequently removing the bran.

10 Taking durum wheat as example, the caryopses have a central nucleus consisting of endosperm, a germ, and a coating of various layers constituting the bran and representing about 15% by weight of the caryopsis itself.

15 The traditional process for the production of flour or semolina foresees a plurality of conditioning phases (3-4) of the caryopses with water and/or steam for 8-12 hours, with the purpose of giving uniformity to and optimising the moisture of the endosperm and at the same time facilitating the separation of the bran. This is followed by milling (breaking, sifting) done by machines provided for that purpose.

20 The finest fractions, consisting mainly of endosperm, together with bran and germ, and separated out by sifting are sent on to successive purification phases.

The coarser fraction, consisting mainly of bran but also of endosperm, is sent to a further milling phase to recover the endosperm.

25 This can be repeated many times, each time producing fine bran and germ particles that are difficult to separate from the endosperm.

This way the final yield of semolina is penalised, as is the quality of the obtained semolina.

Patent EP 0 598 022 describes a procedure for the wetting of cereal grain

- 2 -

that utilises a machine capable of generating vibrations and that leads to a substantial reduction of the time of the single successive conditioning step, with a positive effect on the overall time of the production process.

5 In patent application EP 0 295 774 there is proposed, for the production of wheat flour, an alternative process that mills the caryopses after the selective removal of the various layers constituting the bran.

More precisely, the process according to the aforesaid application provides for the steps of

- 10 1) treating the caryopses with a quantity of water sufficient to condition the outer layers of bran but maintaining the endosperm essentially protected from the water;
- 2) subjecting the thus treated caryopses to friction operations to remove the outermost layers of bran and to successive operations of friction and abrasion to eliminate most of the remaining bran;
- 15 3) conditioning the caryopses with water in a conventional manner and grinding them.

In this way the quantity of bran produced in the milling step is slightly less and the number of crushing steps can be reduced, improving plant productivity.

20 According to the teaching of the abovementioned patent application, it is important that the initial conditioning step of the caryopses be carried out with an amount of water of less than 2% of the caryopses weight to avoid the different bran layers fusing together and those more internal remaining attached to the endosperm, making their separation from this
25 last more difficult.

Nevertheless the Applicant has found experimentally that the process according to this document of the prior art is certainly appropriate for obtaining different bran layers in good purity through the operations of friction and abrasion described in the application but that it leads to
30 serious inconveniences with regard to the subsequent milling phase, especially when working on the industrial scale.

- 3 -

Indeed, the conventional conditioning phase (with a water amount around 16% by weight of the caryopses weight), carried out on the caryopses stripped of most of their bran husk, gives rise to a packing of the caryopses, these tending to adhere one to the other, with consequent insurmountable technical difficulties in their handling and feeding into the milling machines.

The problem underlying the present invention has therefore been one of making available a process for the production of wheat flour or semolina, in particular the semolina of durum wheat, that permits improvement in the yield of the milling phase with the contemporary reduction of production times and the obtaining of a semolina that adheres perfectly to the severe qualitative regulations in force in Italy. All this avoiding the inconveniences indicated further above in relation to the process according to EP 0 295 774.

15 Summary of the invention

Such a problem is solved, according to the invention, by a process for the production of flour or semolina, from soft wheat or durum wheat respectively, starting from the relative caryopses as such, that includes the steps of

- 20 a) wetting said caryopses with such a quantity of water to bring their moisture content to at least 15%, subjecting the caryopses to intense vibrations;
- b) subjecting the wet caryopses to a conditioning step;
- c) subjecting the conditioned caryopses to decortication operations, consisting essentially of an abrasion to remove the outer bran layers;
- 25 d) milling the conditioned and decorticated caryopses.

Preferably the caryopses undergo vibration at frequencies between 50 and 300 Hz, advantageously 75 Hz, generated by means of mechanical, electrical or magnetic energy or else by ultrasound.

Advantageously, the caryopses are exposed to vibration by means of a

- 4 -

machine produced by the company Gräf GmbH & Co. KG, Lahnau, Germany, marketed by the name "Vibronet®.

The conditioning step carried out after the caryopses had been subjected to intense vibrations is only one and is advantageously faster with respect to the multiple (3-4) conditioning steps done conventionally, being limited to 4-6 hours instead of the total of 8-12 hours foreseen by traditional conditioning.

The process according to the present invention has the great advantage, with respect to that traditionally employed, that does not foresee the decortication of the caryopses before the milling step, to attain a 15- 20 % increase in the hourly throughput of one and the same mill. At the same time, it avoids the inconveniences tied to the presence of bran, like for example flooding and blocking due to variations in the volume of the bran as a function of its moisture content.

Moreover the lower amount of bran and germ in the caryopses undergoing milling leads to the production of a flour and semolina with an ash content within the norm even with a greater recovery of the finest fractions with high ash content, with evident advantages for what concerns the semolina and flour yield.

With respect to the process according to EP 0 295 774, the process of the present invention gives the marked advantage of avoiding the packing of the caryopses stripped of their bran husk, that occurs during the wetting and conditioning step of the above patent, and still allows easy decortication of the caryopses.

In fact, by using the process according to the invention, the fusion of the endosperm with the more internal bran layers complained of in the EP 0 295 774 application does not occur. According to a non-binding hypothesis, it is felt that this depends on the fact that the vibrations imparted to the caryopses during the wetting phase lower the surface tension of the bran layers and allow the water to penetrate rapidly to the central part of the caryopsis and become concentrated in the endosperm, leaving the bran layers relatively dry. In this way fusion between the endosperm and the innermost bran layers is avoided.

- 5 -

Furthermore it was found that the friction operations described in the application EP 0 295 774 can be omitted without in any way compromising the yield of the overall process or the quality of the final flour. The removal of the bran from the caryopses is therefore entrusted,
5 according to the present process, solely to abrasion operations, with evident advantages in terms of plant simplification and reduction of processing times.

The process according to the present invention will be further described making reference to an example supplied here in the following as
10 illustrative and not limiting.

EXAMPLE

115 kg of durum wheat previously cleaned conventionally had such a quantity of water added to it as to bring the moisture content of the caryopses to 16.5% and was, at the same time, exposed to vibratory stress
15 imparted by a Vibronet® machine (vibratory pulses lasting about 10 seconds and with a frequency of 75 hz). After a single conditioning of 5 hours, the caryopses were fed to a first decortication or abrasion machine, comprising a rotating shaft with vertical axis supporting abrasive Carborundum wheels that peel away the outermost bran layers
20 in the form of bran powder recovered by suction. The caryopses leaving the first decortication/abrasion machine underwent two successive steps in another two decortication machines that provided respectively for the elimination of the intermediate and internal bran layers, always in the form of bran powder. The three bran powders obtained at the outlet of the
25 three decortication machines differ in fibre content, that was maximum in the first bran powder and minimum in the third, and in protein content, this, vice versa, being maximum in the third bran powder and minimum in the first

The caryopses leaving the third decortication machine, of total weight
30 about 100kg corresponding to 87% of the initial weight, underwent air-stream dynamic cooling before being sent to a conventional mill for grinding. Here the decorticated caryopses were ground according to the operative modalities usually used for milling caryopses as such, obtaining at the end 85 kg of durum wheat semolina (74% yield) fully conforming to

- 6 -

qualitative law requirements. In particular, the ash content was found to conform to the standards as did the particle size and the so-called speckling (in Italian: "puntatura") that, indeed, decreased in both its bran and black components with respect to that of semolina obtained using conventional processes.

Finally it is to be noted that a comparative test carried out on 100 kg of durum wheat from the same lot, subjected to conventional wetting and conditioning for 10 hours with such a quantity of water as to bring the moisture content of the wheat to 16.5%, and then ground without any prior decortication, in the same experimental and operative conditions as the above example, gave 70 kg of semolina, a yield equal to 70%.

As is noted on comparing the two tests, the process according to the invention supplies a higher yield than that of the conventional process. But there is another important advantage of the process according to the invention, and this consists of the possibility of notably increasing the plant milling capacity because the caryopses bear a very reduced bran residue amount and there is no further need to carry out any of the numerous passages (breaking) typical of the conventional milling process.

In the course of the operations on the industrial scale it was confirmed that the flow rate of the decorticated wheat fed to the mill was 15.20 t/h against a flow rate of 12.87 t/h found in the same mill when it was fed non-decorticated wheat.

Always operating on the industrial scale, the process according to the invention resulted in a 75.2 % yield of semolina by weight of the initial durum wheat, whereas the maximum yield obtained with the conventional processes in one and the same mill was 70.1%.

- 7 -

CLAIMS

1. A process for the production of wheat flour or semolina starting from respective caryopses as such, that includes the steps of
 - a) wetting said caryopses with such a quantity of water as to bring their moisture content to at least 15%, by subjecting them to intense vibration;
 - b) subjecting the wet caryopses to a conditioning step;
 - c) subjecting the conditioned caryopses to decortication operations, consisting essentially of abrasion, to remove the bran;
 - 10 d) grinding the conditioned and decorticated caryopses.
2. A process according to claim 1, in which the said wetting step is carried out while subjecting said caryopses to vibrations of a frequency between 50 and 300 Hz.
3. A process according to claim 2, in which said vibrations are generated
15 by means of mechanical, electrical or magnetic energy or else by ultrasound.
4. A process according to claim 3, in which said conditioning step has a duration of 4-6 hours.
5. A process according to any one of the preceding claims, in which said
20 caryopses as such are caryopses of durum wheat.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/10804

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 B02B1/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 B02B B02C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 3 399 839 A (HERMAN ANDERSON ET AL) 3 September 1968 (1968-09-03) column 1, line 9 - line 12 column 1, line 68 - line 71 column 2, line 48 - column 4, line 8 figure 1	1-5
Y	US 5 586 492 A (GRAEF DIETER O) 24 December 1996 (1996-12-24) column 1, line 62 - column 2, line 55 column 3, line 1 - line 4	1-3
Y	US 4 189 503 A (GIGUERE JAMES R) 19 February 1980 (1980-02-19) column 3, line 56 - line 66 column 6, line 52 - line 61 column 9, line 30 - line 46	1,4
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

29 January 2004

Date of mailing of the international search report

09/02/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Redelsperger, C

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/10804

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 4 986 997 A (POSNER ELIESER S ET AL) 22 January 1991 (1991-01-22) column 4, line 59 column 5, line 7 - line 28 -----	1,5

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Publication No

PCT/EP 03/10804

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 3399839	A	03-09-1968	NONE	
US 5586492	A	24-12-1996	DE 4127290 A1 AT 122933 T AU 2406492 A BR 9203191 A CA 2115832 A1 CZ 9400324 A3 DE 4292660 D2 DE 9110617 U1 DE 59202365 D1 WO 9303843 A1 EP 0598022 A1 ES 2072764 T3 HK 41397 A JP 2646171 B2 JP 6213678 A JP 8022386 B JP 6504722 T US 5512310 A	18-02-1993 15-06-1995 16-03-1993 06-04-1993 04-03-1993 19-10-1994 21-12-1995 07-11-1991 29-06-1995 04-03-1993 25-05-1994 16-07-1995 11-04-1997 25-08-1997 05-08-1994 06-03-1996 02-06-1994 30-04-1996
US 4189503	A	19-02-1980	US 4301183 A US 5250313 A	17-11-1981 05-10-1993
US 4986997	A	22-01-1991	NONE	